

MATERIAL DATA SHEET**Sumo-1 (SMT3C), human recombinant**
Cat. # UL-712

The ubiquitin-like SUMO-1 is conjugated to a variety of proteins in the presence of UbcH9 and the SAE1/SAE2 (human) or Aos1/Uba2 (yeast) activating enzyme. SUMO-1 is derived from the precursor pro-SUMO-1 (Accession # NM_003352). Human SUMO-1 shares 46% and 47% identity with SUMO-2 and SUMO-3 respectively. SUMOylation can occur without the requirement of a specific E3 ligase activity, where SUMO is transferred directly from UbcH9 to specific substrates. SUMOylated substrates are primarily localized to the nucleus (RanGAP-1, RANBP2, PML, p53, Sp100, HIPK2) but there are also cytosolic substrates (I κ B α , GLUT1, GLUT4). SUMO modification has been implicated in functions such as nuclear transport, chromosome segregation and transcriptional regulation.

Product Information

Quantity:	500 μ g
Stock:	X mg/ml (X μ M) in 50 mM HEPES pH 8.0, 150 mM NaCl, 1mM DTT. Actual protein concentration will vary with specific Lot #.
MW:	11.1 kDa
Purity:	> 95% by SDS-PAGE

Use & Storage

Use:	Typical <i>in vitro</i> concentrations for conjugate formation is 10-50 μ M depending on conditions.
Storage:	Store in -80°C. Avoid multiple freeze/thaw cycles.

Literature

References:	Desterro J.M., <i>et al.</i> (1997) <u>FEBs. Lett.</u> 417 :297-300 Okama T., <i>et al.</i> (1999) <u>Biochem. Biophys. Res. Comm.</u> 254 :693-698 Seeler J-S. and Dejean A. (2003) <u>Nat. Rev.</u> 4 :690-699 Su H-L., <i>et al.</i> (2002) <u>Gene</u> 296 :65-73 Tatham M.H., <i>et al.</i> (2001) <u>J. Biol. Chem.</u> 276 :35368-35374 Yeh E.T.H., <i>et al.</i> (2000) <u>Gene</u> 248 :1-14
--------------------	--

For Laboratory Research Use Only, Not For Use in Humans