

MATERIAL DATA SHEET**20S Proteasome, mouse****Cat. # E-355**

The 20S Proteasome is the catalytic core component of the multi-complex 26S Proteasome that selectively degrades intracellular proteins. It is commonly associated with regulatory complexes, which include the 19S Proteasome, the PA28 alpha/beta complex, or the PA28 gamma complex. The 20S Proteasome is composed of 28 subunits arranged into four stacked rings. The outer rings, containing seven subunits each, are composed of closely-related but non-identical alpha subunits. The amino-terminal tails of the alpha subunits form a gate that restricts substrate entry into the catalytic core. The inner rings, also containing seven subunits each, are composed of closely-related but non-identical beta subunits. The amino-terminal tails of six of the beta subunits, three per ring, have proteolytic activity. Inhibition of 20S Proteasome proteolytic core activity using small molecule inhibitors is a valuable tool for the functional study of a variety of proteins and for therapeutic intervention. The 20S Proteasome can be activated chemically by the addition of detergent, by the proteinaceous activator PA28 Activator alpha (E-380), or by 19S/PA700 (E-366).

Product Information

Quantity:	50 µg
Stock:	X mg/ml (X µM) in 50 mM HEPES pH 8, 150mM NaCl, 1mM DTT.
MW:	700 kDa
Purity:	> 95% by SDS-PAGE under reducing conditions and visualized by Colloidal Coomassie Blue stain.

Use & Storage

Use:	The Murine 20S Proteasome is able to degrade substrates in an ATP-independent manner. It can be activated chemically with SDS (0.035%) or by the addition of PA28. Reaction conditions will need to be optimized for each specific application. We recommend an initial 20S Proteasome concentration of 0.5-5 nM.
Storage:	Store at -80°C. Avoid multiple freeze/thaw cycles.

Literature

References:	Stadtmueller, B.M. & C.P. Hill (2011) <u>Mol. Cell</u> 41 : 8 Kim, H.M. <i>et al.</i> (2011) <u>Biochim. Biophys. Acta</u> 1809 : 67 Xie, Y. (2010) <u>J. Mol. Cell Biol.</u> 2 : 308 Kisselev, A.F. <i>et al.</i> (2012) <u>Chem. Biol.</u> 19 : 99 Ma, C.P. <i>et al.</i> (1992) <u>J. Biol. Chem.</u> 267 : 10515□
--------------------	--

For Laboratory Research Use Only, Not For Use in Humans

Rev: 02/04/2014

840 Memorial Drive, Cambridge, MA 02139 Phone: 617-576-2210 FAX: 617-492-3565
www.bostonbiochem.com

The contents of this datasheet (unless otherwise noted) are Copyright © 2008 Boston Biochem, Inc. All rights reserved. Duplication in whole or in part is strictly prohibited without the express written consent of Boston Biochem, Inc. "Boston Biochem" is a Trademark of Boston Biochem, Inc., 840 Memorial Drive, Cambridge, MA 02139.