

MATERIAL DATA SHEET

3CL Protease, SARS-CoV-2 virus recombinant Cat. # E-720

The 3CL protease (aka 3CL^{pro}, M^{pro} or "Main Protease) from the human SARS-CoV-2 coronavirus (Severe Acute Respiratory Syndrome coronavirus 2) is a C30-type cysteine protease located within the non-structural proteins 3 (NS3) section of the viral polypeptide. 3CL^{pro} activity is required to process the viral polyprotein into functional, mature subunits, and there are 11 or more sites of cleavage, many containing the sequence LQ[S/A/G]; the protease cleaves c-terminal to the glutamine amino acid. Along with the CoV-2 Papain-Like Protease (catalog number E-611), 3CL^{pro} presents an attractive target for therapeutic intervention for COVID-19. Because no human proteases with a similar cleavage specificity are known, inhibitors of 3CL^{pro} are unlikely to cause mechanism-based toxicity. This recombinant protein is untagged.

Product Information

Quantity: 50 μg

Stock: 1.7 mg/ml (50 μM) in 50 mM HEPES pH 7.5, 150 mM NaCl, 1 mM DTT,

1 mM EDTA.

MW: 34 kDa

Purity: > 95% by SDS-PAGE under reducing conditions and visualized by Colloidal

Coomassie Blue stain

Use & Storage

Recombinant SARS-CoV-2 virus 3CL Protease is a cysteine protease that cleaves proteins with sequences including a LQ\[S/A/G]. Reaction conditions

will need to be optimized for each specific application.

Storage: Store at -80°C. Avoid multiple freeze/thaw cycles.

Literature

References: Chen Y.W. et al. (2020) <u>F1000Research</u> doi: 10.12688/f1000research.22457.2

Pillaiyar T., et al. (2016) J. Med. Chem. **59**: 6595

Zhang L. et al. (2020) Science doi: 10.1126/science.abb3405

For Laboratory Research Use Only, Not For Use in Humans

Rev: 5/20/2020

840 Memorial Drive, Cambridge, MA 02139 Phone: 617-576-2210 FAX: 617-492-3565 www.bostonbiochem.com