
MATERIAL DATA SHEET

SUMO2 Biotin, *human recombinant***Cat. # UL-754**

SUMO-2 modified with biotin via primary amine coupling resulting in modification of lysine residues as well as the N-terminus. Although having a fully functional C-terminus, lysine modification may limit the ability of this reagent to propagate poly-SUMO chains. Biotinylated SUMO-2 can be detected using avidin-linked reagents for higher efficiency and sensitivity than with antibodies.

Product Information

Quantity:	50 µg
Stock:	X mg/ml (XµM) in 50mM Hepes pH 7.5, 100mM NaCl. Actual concentration varies with lot number.
MW:	10.6 kDa
Purity:	95% by PAGE

Use & Storage

Use:	Depending on experimental conditions and detection method, suggested <i>in vitro</i> concentrations for thiolester formation is 5 to 20 µM.
Storage:	Store at -80°C. Avoid multiple freeze/thaw cycles.

Literature

References:	Desterro, J.M. <i>et al.</i> (1997) <u>FEBs. Lett.</u> 417 :297-300 Okama T. <i>et al.</i> (1999) <u>Biochem. Biophys. Res. Comm.</u> 254 :693-698 Seeler J-S. and Dejean A. (2003) <u>Nat. Rev.</u> 4 :690-699 Su H-L. <i>et al.</i> (2002) <u>Gene</u> 296 :65-73 Tatham M.H. <i>et al.</i> (2001) <u>J. Biol. Chem.</u> 276 :35368-35374 Yeh E.T.H. <i>et al.</i> (2000) <u>Gene</u> 248 :1-14
--------------------	--

For Laboratory Research Use Only, Not For Use in Humans