

MATERIAL DATA SHEET**Ubiquitin Mutant G76A, human recombinant****Cat. # UM-G76A**

Mature Ub has a highly conserved C-terminal diglycine motif which is crucial for activity and recognition with conjugation and deconjugation enzyme components. The replacement the last glycine residue with alanine results in a Ub that supports E1-Ub thioester formation and downstream conjugation reactions (transfer to E2, E3) but at a rate ~20% compared to wildtype Ub. This mutant however, inhibits deconjugation and prevents the removal of Ub from modified protein substrates by deubiquitinating enzyme (DUBs). Since this Ub becomes irreversibly conjugated to protein, it shifts the equilibrium between the bound and unbound form in the direction of conjugation, at the expense of the free form.

Product Information

Quantity:	100 µg, lyophilized powder
Solubility:	Aqueous solutions up to 5 mg/ml
Purity:	> 95% by SDS-PAGE
MW:	8.6 kDa

Use & Storage

Use:	Typical concentrations will depend on specific assay conditions and method of detection.
Storage:	Solubilized solution at -20°C. Avoid multiple freeze/thaw cycles.

Literature

References:	Dayal S. <i>et al.</i> (2009) <u>J.Biol.Chem.</u> 284 :5030-5041
	Geng F. and Tansey W.P. (2008) <u>Mol. Biol. Cell.</u> 19 :3616-3624
	Pickart C.M. <i>et al.</i> (1994) <u>J.Biol.Chem.</u> 269 :7115-7123
	Ravid T. and Hochstrasser M. (2007) <u>Nat. Cell Biol.</u> 9 :422-427
	Wilkinson K.D. <i>et al.</i> (1995) <u>Biochem.</u> 34 :14535-14546

For Laboratory Research Use Only, Not For Use in Humans

840 Memorial Drive, Cambridge, MA 02139 Phone: 617-241-7072 FAX: 617-492-3565

www.bostonbiochem.com

The contents of this datasheet (unless otherwise noted) are Copyright © 2008 Boston Biochem, Inc. All rights reserved. Duplication in whole or in part is strictly prohibited without the express written consent of Boston Biochem, Inc. "Boston Biochem" is a Trademark of Boston Biochem, Inc., 840 Memorial Drive, Cambridge, MA 02139